



ESTRO technical guideline: intensity modulated radiotherapy and image guided radiotherapy for rectal cancer

ARTICLE INFO

Keywords:

Rectal cancer
Intensity modulated radiotherapy
Volumetric modulated arc therapy
Image guided radiotherapy
Guidelines

ABSTRACT

Introduction: Intensity Modulated Radiotherapy (IMRT) and Image Guided Radiotherapy (IGRT) have become an integral part of standard care for rectal cancer, but evidence-based detailed guidance is lacking to support its clinical implementation and use. This European Society for Radiotherapy and Oncology (ESTRO) technical guideline aimed to assess the available evidence and provide recommendations for their use in rectal cancer treatment.

Materials and methods: The ESTRO Lower Gastrointestinal (GI) Cancer guidelines subcommittee formed a writing panel to address key questions (KQs) on the application of IMRT and IGRT in rectal cancer management. The panel conducted a literature review and where evidence was insufficient, expert consensus was used.

Results: The writing panel (2 RTTs, 2 medical physics experts, 3 radiation oncologists) identified 14 KQs. Recommendations were based on low to moderate evidence and/or expert consensus. Supine positioning is preferred for patient comfort and stability. Comfortable full bladder should be used. For OAR delineation and dose optimization, mandatory (Bladder, Bowel Cavity, Femoral Heads) and optional OAR were defined. Auto-delineation is supported for OAR and can be considered for target volumes. Dose metrics for minimizing gastrointestinal and genitourinary toxicity were identified. Auto-planning tools and MR-only workflows are feasible, both require proper QA. Uncertainties from setup and target volume shape variations must be accounted for with appropriately defined margins. Daily volumetric image guidance is recommended for treatment verification.

Conclusion: This ESTRO technical guideline for the use of IMRT and IGRT for rectal cancer was developed to support development and implementation into clinics.

Introduction

The standard treatment strategy for many rectal cancer patients includes the use of radiotherapy, either short course radiotherapy (SCRT, 25 Gy in 5 fractions) or long course radiotherapy (LCRT, 45–50.4 Gy in 25–28 fractions, 1.8–2.0 Gy per fraction, with optional tumour boost) with concomitant fluorouracil-based chemotherapy. This will most commonly be in the pre-operative setting, where it improves local control in cases of intermediate risk, resectable cancer and facilitates downstaging in high-risk borderline resectable or initially unresectable tumours [1,2]. However, (chemo)radiotherapy is also increasingly being used to facilitate non-operative management [3–5] of early rectal cancers [6,7]. This may allow patients to avoid the substantial morbidity and potential diminished quality of life associated with surgery, but may impose greater demands on the quality of radiotherapy.

Two key techniques are essential for modern radiotherapy planning and delivery: (1) Intensity modulated radiotherapy (IMRT) or volumetric modulated arc therapy (VMAT), which allow for precise dose shaping using steep dose gradients around the clinical target volume (CTV), and (2) image guided radiotherapy (IGRT (2D/3D)), which visualizes patients anatomy (2D or 3D, bony anatomy and/or soft tissue) and thus improves treatment accuracy. Combining IMRT and IGRT enables optimized planning target volume (PTV) margins. These techniques have the potential to significantly reduce radiation exposure to healthy tissues, particularly bladder and bowel, without compromising target volume

coverage. Retrospective studies have indicated that IMRT may reduce acute toxicity for rectal cancer patients undergoing preoperative (chemo) radiotherapy [8], although the picture has been nuanced by more recent publications [9]. IMRT and IGRT combined facilitate the delivery of simultaneous integrated boosts (SIB) [10–12]. Both techniques are currently widely available, and are part of standard treatment strategies for rectal cancer radiotherapy in many countries [11,13]. A recent UK survey reported 95% of centres treated all patients with IMRT [14].

However, there are limited practical and technical international guidelines to support the implementation of IMRT and IGRT for rectal cancer. Some national guidance exists [1,11,13] but there is a lack of published evidence-based guidelines. Therefore, the European Society for Radiotherapy and Oncology (ESTRO) and its Lower GI guidelines subcommittee tasked an initiative to develop a technical guideline to support the clinical application of IMRT and IGRT. This manuscript presents the technical guideline, reviewing the available literature and providing recommendations for effective implementation to maximize clinical benefits.

Materials and methods

This guideline, along with its scope and the composition of the writing and review panel, were approved by the ESTRO Guidelines Committee. The composition of the writing panel aimed for diversity in discipline, experience and a balance between academic and non-

<https://doi.org/10.1016/j.radonc.2026.111409>

Received 28 January 2026; Accepted 28 January 2026

Available online 7 February 2026

0167-8140/© 2026 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

academic radiotherapy centres. The writing panel comprised an international group of experts, including radiation oncologists (RO) (n = 3), medical physics experts (MPE) (n = 2), and radiation therapists (RTT) (n = 2). The panel was chaired by AA and RdJ.

The scope and content of this guideline has been structured around a series of Key Questions (KQs). KQs were developed based on structured discussions within the writing group during online meetings, focused on the radiotherapy workflow and with particular emphasis on IMRT and IGRT. Resulting KQs were reviewed and approved by the Lower GI guidelines committee (ESTRO). Clinical and Gross target volume definitions were excluded [15]. Adaptive radiotherapy, such as library of plans or online adaptive radiotherapy, were also excluded on the basis that these techniques and technologies are not (yet) widely available.

A comprehensive literature review (PubMed only) was conducted in accordance with PRISMA guidelines to address the KQs. A standardized set of search terms and inclusion/exclusion criteria were applied across all KQs to identify papers on rectal cancer and (chemo)radiotherapy. General inclusion criteria allowed for both systematic reviews and original research articles and studies conducted in both the preoperative and non-operative management setting. Additional search terms were specific to each KQ; see Supplement A for full details. The KQ-specific literature searches were performed in PubMed between November 2023 and June 2025. Supplement A details whether abstracts and/or grey literature were considered for each KQ.

The findings from the literature reviews served as the basis for the recommendations for each KQ. The quality of evidence and strength of recommendations were assessed following the methodology established by the American Society for Radiation Oncology (ASTRO) (ASTRO Methodology Guide – V1: 5/2019). In cases where the available literature was insufficient, expert consensus was achieved through discussion and/or voting. Consensus was defined as at least a majority agreement (n ≥ 4). When the group was unanimous in its opinion, we labelled the strength of recommendation as ‘strong’.

All members of the writing group contributed to the final manuscript, which underwent peer review by an independent reading panel. The final version was approved by all members of the writing group.

Results

The writing panel identified 14 KQs. The findings from these KQs (based on literature review and/or expert consensus) were synthesized into recommendations, which are summarized in Table 1, as well as visualized in a Flowchart (Fig. 1).

KQ 1) What is the optimal patient immobilization with respect to reproducibility, stability and dose to organs at risk?

The prone position with a bellyboard device was commonly used during the era of 2D and simple 3D treatment planning (three or four field box techniques) to reduce dose to normal tissue by displacing the small bowel from the target volume [16–19], with or without bladder distension [20].

The transition to IMRT / VMAT and IGRT, and the associated optimisation of plan conformality and PTV margins, has allowed for a reduction in planned bowel dose. Studies have investigated whether the prone position with a bellyboard device is still beneficial in this context. Most studies have found that it can reduce radiation dose to the small bowel further [21–29], but not all publications support this conclusion [30,31]. Bladder filling / distension also appear to maintain some benefit in this setting [26,28]. However, these results are based on relatively small populations and on comparison of treatment planning that does not take into account inter- and intrafraction motion (both setup variation and soft tissue changes).

Treatment position also impacts setup reproducibility and stability. Reproducibility appears superior in the supine position relative to the prone position (with or without bellyboard) [22,26,29,30,32]. Literature on intrafraction motion is limited, but the supine position is also favoured with respect to stability [33]. Finally, the supine position is

Table 1
Overview of key questions and recommendations.

	Key Questions	Recommendations	Strength	Level of Evidence
1.	What is the optimal patient positioning with respect to reproducibility, stability and dose to OAR?	• The supine position should be the first option for treatment position to enhance patient comfort, compliance and stability. • For individual patients where reduction of bowel dose is the primary concern, prone position on bellyboard should be an alternative option.	Strong	Low/ expert consensus
2.	What is the optimal regimen of patient preparation (bowel and bladder)?	• A patient preparation protocol should be used for a comfortable full bladder for planning and treatment (300–500 ml, 30–45 prior to imaging/treatment). Patients should be educated on maintaining hydration status for both planning and treatment. • Strategies should be used to minimise flatus/faeces in the rectum for planning and treatment. An assessment of rectal diameter should be performed to ensure the CT planning scan is representative of patient anatomy.	Strong	Low/ expert consensus
3.	What are the mandatory and optional OAR to be outlined to support treatment planning and evaluation?	Mandatory OAR: • Bladder • Bowel Cavity • Femoral Heads Recommended OAR: • Vagina • External Genitalia • Penile Bulb • Anal Sphincter • Sacral Bones • Stoma (when within 1 cm of PTV in cranio-caudal direction)	Strong	Expert consensus
4.	Which consensus guidelines are there on the definition of OARs to support their outlining?	• Consensus guidelines for nomenclature and outlining of OAR should be used to enhance safety and improve quality for clinical practice as well as to enable data pooling. For standardized nomenclature, AAPM TG-263 is recommended; for outlining of OARs, see Table 2 for recommendations.	Strong	Expert consensus
5.	Can auto-delineation be used	• Auto-delineation can be used for OAR segmentation, with	Conditional	Low/ expert consensus

(continued on next page)

Table 1 (continued)

	Key Questions	Recommendations	Strength	Level of Evidence
	for target volumes and/or OAR?	significant time-reduction benefits. • Auto-delineation can be considered for target volume segmentation. • When implementing consider both qualitative acceptability criteria (e.g. geometric accuracy, dosimetric impact) and outcomes assessments (e.g. actions to be taken for corrections); with subsequent use dependent on both case-specific QA and routine model QA.		
6.	What are the mandatory and optimal dose levels and metrics to consider for OAR, in order to minimize both acute and late toxicity?	• Dose to small bowel loops or bowel cavity should be limited to prevent acute and late GI toxicity. A range of optimisation objectives covering low to high dose levels are recommended. • The bowel cavity, or a comparable volume (such as peritoneal space), should be used for plan optimization, irrespective of the volume used for plan evaluation and reporting. • Dose to bladder should be limited, with a particular focus on limiting $V_{30-35Gy}$ to prevent acute toxicity. • For any other OAR outlined for an individual patient, dose should be reviewed as part of plan evaluation and prescription, and any patient-specific considerations should be recorded.	Strong	Low/expert consensus
7.	Which dose metrics are to be considered for target dose evaluation?	• Radiotherapy treatment planning should aim to cover the planning target volume for primary tumour, involved lymph nodes, and elective volumes with 95% of the prescription dose, while minimising hotspots. • If delivering a SIB to tumour or involved nodes, this should be as conformal as possible, minimising spill-over of high	Strong	Expert consensus

Table 1 (continued)

	Key Questions	Recommendations	Strength	Level of Evidence
8.	How should dose to OAR and target volumes be prioritized during treatment plan optimization and evaluation?	• dose into surrounding elective volumes. • In general, dose to the target volume should take precedence over dose to OAR, although this prioritization may be adjusted for individual patients. • When prioritization changes, the applied registration strategy for daily IGRT should align accordingly.	Strong	Expert consensus
9.	Are there any specific technical considerations to be made regarding IMRT planning?	• Guidance provided separately in the main text and Table 4.	—	—
10.	Can auto planning solutions add value to IMRT planning, and what is the required QA?	• Commercially available auto-planning solutions can be used to improve the OAR sparing and increase the time efficiency and consistency of IMRT planning. • When used, verify and monitor local clinical performance continuously.	Conditional	Low
11.	What is the optimal method to calculate and report dose?	• Dose should be calculated and reported in dose-to-medium (D_m) where possible. • Special care should be taken when comparing D_m with dose-to-water (D_w) in cortical bone. It is not recommended to convert D_m to D_w (dose to water in medium, $D_{w,m}$) as this will result in poorer agreement with either D_w and D_m.	Strong	Expert consensus
12.	Is an MRI only workflow feasible using (commercially available) sCT for treatment planning?	• Commercially available sCT generation software can be used to generate sCT for dose calculation in an MR-only workflow. • MRI or sCT can be used as reference image for position verification in an MR-only workflow for rectal cancer; but additional QA may be needed due to the lack of native support in commercial software.	Conditional	Moderate/expert consensus
13.	How are large are setup and target volume shape uncertainties?	• Systematic and random errors in setup and target volume shape variation are	Strong	Moderate

(continued on next page)

Table 1 (continued)

Key Questions	Recommendations	Strength	Level of Evidence
14. What is the optimal IGRT protocol to verify dose delivery?	substantial, even with daily image guidance, and should be accounted for in target volume margins. • The required margin should be defined separately for the primary tumour, involved lymph nodes, mesorectum (upper and lower) and further elective volumes (extra-fascial lymph nodes). • Daily volumetric imaging (i.e., Conebeam-CT/3D-MV) should be used to correct systematic and random setup errors by matching the bony structures as well as to evaluate the position of target volumes and OAR. • Supporting protocols (e.g., traffic light) should be used for image evaluation and decision making.	Strong	Moderate/expert consensus

Recommendations are based on evaluation of multiple factors including the quality of evidence (QoE), individual study quality, and expert consensus, all of which inform the strength of recommendation. QoE is based on the body of evidence available for a particular key question and includes consideration of number of studies, study design, adequacy of sample sizes, consistency of findings across studies, and generalizability of samples, settings, and treatments (ASTRO methodology guide). In cases where the available literature was insufficient, expert consensus was achieved through discussion and/or voting. Consensus was defined as at least a majority agreement ($n \geq 4$). When the group was unanimous in its opinion, we labelled the strength of recommendation as ‘strong’.

Abbreviations: Image guided Radiotherapy (IGRT); Intensity Modulated Radiotherapy (IMRT); Organ(s) at Risk (OAR); Quality Assurance (QA); Gastro-intestinal (GI); Magnetic Resonance Imaging (MRI); Synthetic Computed Tomography (sCT); Simultaneous Integrated Boost (SIB).

documented as superior in terms of patient comfort; especially for frail patients, post-surgical cases, and patients with stomas [22,27,30].

The expert consensus is unanimously to use the supine position as the first choice prioritizing factors such as interfraction setup reproducibility, intrafraction stability as well as patient comfort. However, if minimizing small bowel dose is the only or primary objective the prone position with a bellyboard device is still optimal when using IMRT/VMAT.

KQ 2) What is the optimal regimen of patient preparation (bowel and bladder)?

Bowel and bladder filling can change considerably between treatment planning and delivery, and between individual treatment fractions [34,35]. However, there are limited data to support a specific bowel or bladder preparation protocol.

Theoretically, a (comfortable) full bladder pushes small bowel away from the target volume, thus minimising dose to the anterior part of the bladder and small bowel. To enhance bladder volume consistency throughout radiotherapy, it is recommended (expert consensus, unanimously) to provide patients with structured education and clear drinking instructions to ensure optimal hydration and a comfortable full bladder during both planning and treatment sessions (300–500 ml,

30–45 min prior)[11]. However, factors such as chemotherapy or medications may affect hydration status and thus bladder filling. Ultrasound scanners to assess bladder volume are very accurate and can assist in optimizing bladder volume just prior to treatment but may be logistically challenging [36,37]. Volumetric imaging is useful for anatomical assessment and volume prediction [38]. Some studies describe bladder volume control via catheter irrigation [39,40], and bladder volumes can decrease over the course of treatment [36,37,40].

Patients should be advised to empty their bowels before planning and treatment to reduce target volume deformation and unnecessary organ(s) at risk (OAR) dose. The rectum should not be extended (e.g. diameter should not exceed 4 cm [11]) at the planning stage to ensure that planning imaging is representative of reproducible anatomy throughout the duration of treatment (expert consensus, unanimously). Possible solutions to reduce the variability include allowing patients to pass gas or empty their bowels before proceeding; review of rectum diameter on scout imaging before full scan; review of the rectum diameter on the diagnostic imaging to assess the consistency of the rectal size and filling over time (and hence the feasibility of minimizing the diameter through interventions).

KQ 3) What are mandatory and optional OAR to be outlined to support treatment planning and evaluation?

Outlining of normal tissue as OAR may be for several purposes; including dose plan optimisation and evaluation, reporting of OAR dose for individual patients (even where no clear dose constraints exist), and to support long-term evidence generation. The first two considerations are the main focus of the current guidance.

Due to insufficient and inconsistent guidance in the literature, the writing panel conducted a formal voting process to achieve expert consensus on OAR for delineation (see Supplement B).

The OAR classified as mandatory for delineation were: *Bladder* (Unanimously), *Bowel Cavity* (Majority Agreement), *Femoral Heads* (Unanimously). Several definitions of Bowel Cavity were discussed, including ‘bowel bag’ as defined by the EMBRACE group (peritoneal cavity containing bowel loops, following the outer contour of visible bowel loops) [41], and ‘bowel bag’ from the RTOG atlas (abdominal contents excluding muscle, bones and non-GI normal structures) [42]. The writing panel agreed that the mandatory Bowel Cavity OAR should encompass all anatomy which may potentially contain bowel loops during treatment delivery, in line with the RTOG ‘bowel bag’ definition (see also KQ 6) and ‘Spc.Bowel’ described by Mir et al [43].

The OAR classified as recommended for delineation were (all decided with Majority Agreement): *External Genitalia*, *Vagina*, *Penile Bulb*, *Anal Sphincter*, *Sacral Bones*, and *Stoma* (applicable only when located within 1 cm of the PTV in the cranio-caudal direction).

The OAR classified as optional for delineation were (all decided with Majority Agreement): *Prostate & Seminal Vesicles*, *Ovaries*, *Uterus & Cervix*, *Lumbosacral Plexus*, *Diverticulum* (defined as part of the bowel that ends in the stoma), *Inguinal Area*.

An additional round of voting was conducted to determine the recommendations for OAR delineation in specific clinical scenarios. In these scenarios, the panel aimed to highlight additional OAR that may warrant delineation beyond those classified as mandatory. These were: Intention to boost either the primary tumour (GTV) (*Vagina* and *Anal Sphincter*) or clinically involved lymph nodes (*Sacral Bones* and *Lumbosacral Plexus*), intention for up front organ preservation (*Anal Sphincter*), intention for administering Total Neoadjuvant Treatment (*Bone Marrow*). For an overview see also Table 2.

Finally, the writing panel acknowledges that certain clinical scenarios and factors (such as patient age, frailty, or fertility preservation) can require complex decision making, and covering all scenarios is outside the scope of this current guideline. The panel also identified the rectum wall itself as a potential OAR in the context of organ preservation, but could not provide recommendations on its use due to a lack of literature and clinical experience with plan optimisation and evaluation strategies.




Fig. 1.

KQ 4) Which consensus guidelines are there on the definition of OAR to support their outlining?

A comprehensive understanding of normal tissue dose response is essential for optimal radiotherapy treatment planning. However, there is a significant knowledge gap, particularly regarding IMRT and VMAT, with only limited data available (see also KQ6). One contributing factor to this gap is the variability in delineation of OAR across clinical practice and research studies. Such inconsistencies not only hinder the comparability of dose volume metrics and their correlation with treatment outcomes within and between studies, but also impact clinical outcomes in routine practice.

The adoption of standardized delineation guidelines for normal tissues enhances the consistency and accuracy of radiotherapy, leading to improved clinical outcomes and reduced toxicity [44,45]. Additionally, the implementation of standardized nomenclature as in AAPM TG263

facilitates automation within clinical departments and supports wider research [46]. It is therefore recommended (expert consensus, unanimously) to use consensus guidelines for delineation as well as standardized nomenclature.

A review of existing consensus guidelines was conducted to identify standardized recommendations for outlining OAR in rectal cancer radiotherapy. Key OAR, as defined by expert consensus in KQ3 and their recommended consensus guidelines, are summarized in Table 2 [11,42,43,47–51].

KQ 5) Can auto-segmentation be used for target volumes and/or OAR?

Auto-segmentation is a promising tool for increased contouring consistency and resource savings across radiotherapy [52]. For rectal cancer, literature indicates that auto-segmentation can be used for relevant target volume and OAR segmentation [53–59] with evidence of significant time reduction as a benefit after implementation

Table 2

Overview of organs at risk (OAR) with available guidelines.

OAR	Standardised nomenclature (TG-263)	Reference and description
<i>Mandatory</i>		
Bladder	Bladder	The bladder should be contoured in entirety from base to dome. The lateral extent is the outer bladder wall [42,43].
Femoral heads (left / right)	Femur_Head_R/L	The structure includes the ball of the femur, femoral neck, greater trochanter, and proximal shaft to the caudal limit of the lesser trochanter [42,43].
Bowel cavity	Spc_Bowel*	Contour abdominal contents. Inferiorly from the most inferior small or large bowel loop (excluding rectum). Contour to 2 cm superior to PTV. Exclude: CTV, muscles and major vasculature (common, internal and external iliac vessels) and subtract bladder and uterus (if relevant) from structure.[11,42,43]
<i>Recommended</i>		
Stoma bag	Bag_Ostomy	No consensus guideline available. Should be used as an avoidance structure.
Genitals	Genitals	The female genitals structure encompasses the clitoris, labia major and minora, mons pubis, and the surrounding fat. The male genitals structure encompasses the entire penis, scrotum, perineal body, and the surrounding fat. Both structures extend laterally to the inguinal creases [43,47].
Penile bulb	PenileBulb	The penile bulb is the portion of the bulbous spongiosum of the penis immediately caudal to the genito-urinary diaphragm. The structure is normally 9–10 mm in the cranial-caudal direction. The contour should not continue in to the shaft of the penis [42,43,51].
Sacral bone	Bone_Sacrum*	No consensus guideline available.
Anal sphincter	Sphincter_Anal	No consensus guidelines available. Visuals see Arias et al. [48]. About 1 cm under the last contrast in the rectal ampulla.
Vagina	Vagina	The vagina is a central, muscular tube, approximately 7–9 cm in length, connecting the cervix or vaginal vault to the vulva [50,51].
<i>Optional</i>		
Bone marrow	BoneMarrow	No consensus guideline available as there is no consensus in delineation in the dose–response literature.
Cervix	Cervix	No consensus guideline available.
Diverticulum	Diverticulum*	No consensus guideline available.
Inguinal area	Spc_Inguinal*	Defined as the skin fold. No consensus guideline available. Should be used as an avoidance structure.
Ovaries and fallopian tubes (left / right)	Adnexa_R/L	Ovaries and fallopian tubes [42,50].
Prostate, Seminal vesicles	Prostate; SeminalVesc	Inferiorly from its apex and superiorly to its base. Entire seminal vesicles including those slices that also have prostate identified [42].
Uterus	Uterus	No consensus guideline available.
<i>Where lymph nodes are boosted:</i>		

Table 2 (continued)

OAR	Standardised nomenclature (TG-263)	Reference and description
Lumbosacral Plexus	SacralPlex	The lumbosacral plexus should be contoured from the L4 nerve root to the cranial most portion of the femoral neck [43,49].

*No standardized nomenclature available (American Association of Physicists in Medicine (AAPM) Task Group (TG) report 263); suggested naming provided.

[57,58,60–67]. The systematic literature search did not identify any prospective multicentre clinical implementation studies in the context of radiotherapy for rectal cancer. Existing publications are retrospective in nature and were predominantly conducted in single centre settings, identifying a significant evidence limitation. For most published studies, the number of patients used to train and validate the auto-segmentation models aligned with recommendations [68]. Training sets for these models typically ranged from 50 to 100 patients and testing datasets included at least 20 patients [54–57,59,64,66,67,69–74].

The majority of studies used human observer contours as the ground truth and most reported on in-house models rather than commercial ones. The reasons for this preference were not clear, but may be due to financial reasons, data sharing restrictions or academic incentives. Additionally, commercial systems will contain pre-set models which will have been pre-trained on external datasets, thus not necessarily reflecting local practice.

Published data indicate that for both OAR and target volume, the auto-segmentation solutions evaluated provide acceptable and satisfactory results with considerable time savings [54,55,61–63,65,67,69,75–77]. Additionally, studies evaluating end-users' preferences found that auto-segmentation was preferred in 40–52% of cases versus a human-made segmentation [66,72]. A few authors investigated the target volume or OAR only, while others considered both [54–57,59]. Papers specifically reporting on rectal cancer target volumes evaluated GTV delineation [64,70,72,76], elective CTV volumes [56,58,66,73,77] or primary tumour CTV. However, a large number of publications did not specify the target volume category they were addressing. There were no papers advocating against the use of auto-segmentation tools, either commercial or in-house, as long as structures are being evaluated and adjusted when necessary.

Overall, the limited evidence available supports the use of auto-segmentation for OAR and both elective and primary target volumes. Additionally, the use of auto-segmentation may increase the number of OAR evaluated in routine clinical practice. For implementation of auto-segmentation, qualitative acceptability criteria (e.g. geometric accuracy, dosimetric impact) and outcomes assessments (e.g. actions to be taken for corrections) are needed [60]. Both case specific QA and routine model QA should be used. Clinically implemented models need to be periodically recommissioned, but thresholds are not yet settled in the literature on when to prompt/initiate recommissioning [68].

KQ 6) What are the dose levels and metrics to consider for OAR, in order to minimize both acute and late toxicity?

Understanding normal tissue dose response and dose volume effects is essential for IMRT optimisation and evaluation. Most available data for radiotherapy for rectal cancer are based on 3D-CRT, with limited evidence in the IMRT or VMAT setting. It is unclear if findings from 3D-CRT are generalizable to IMRT, where the DVH correlations from 3D-CRT may be broken. Additionally, data are largely from LCRT, with almost none for SCRT.

The relationship between dose to the small bowel and acute gastrointestinal toxicity during preoperative LCRT has been extensively studied; see [Supplement Table 1](#) and the 2019 review by Holyoake et al [78]. There is evidence to support a correlation between the volume of individual small bowel loops receiving dose levels from 5 Gy to 45 Gy (V_{5Gy} to V_{45Gy}) and acute diarrhoea, with no clear single dose level cut off

identified. Most studies of 3D-CRT found strongest associations with toxicity for low dose volumes ($V_{10-15Gy}$), but this may be an artefact of the statistical analyses [79] and DVH data collinearity. Conversely, limited data from VMAT patients found the strongest association with higher dose volumes (V_{45Gy}) [80]. Similarly, comparative studies of small bowel loops and peritoneal space as OAR with 3D-CRT [81] may not be generalizable to IMRT planning either. Very limited data is available on the association between dose to small bowel loops and late gastrointestinal toxicity [82,83]. A single large study ($n = 486$ patients) found the strongest relationship for V_{30Gy} , and recommended to keep small bowel $V_{30Gy} < 200$ cc (5% risk grade 3 +) or < 300 cc (10% risk grade 3 +). Consequently, optimization objectives should cover a range of dose levels, with priority given to limiting medium to high dose volumes (≥ 30 Gy) to limit late toxicity.

Most available data are based on the outlining of small bowel, being defined as individual bowel loops, rather than a more general ‘bowel cavity’ or ‘peritoneal space’ [81,84]. Given the significant interfractional movement of individual loops [85], the actual dose delivered to OAR may differ from the planned dose [86]. Data from whole pelvis prostate cancer IMRT indicate that treatment plans optimized with a peritoneal space / ‘pelvis cavity’ OAR are significantly more robust to interfractional bowel position variations in terms of the OAR dose delivered. Therefore, the bowel cavity, or a comparable volume, should be used for plan optimization, irrespective of the volume used for plan evaluation and reporting. Dose optimization to individual bowel loops may have to be considered when delivering a high dose boost.

Limited literature has explored the relationship between whole bladder dose and acute cystitis during LCRT (Supplement Table 1). One large retrospective analysis, based on a cohort of patients of which approximately one third were treated with IMRT, suggests that limiting $V_{30-35Gy}$ may reduce acute toxicity [87]. For late toxicity to bladder, there is a lack of data regarding dose volume effects, precluding specific recommendations on dose thresholds. Expert consensus (unanimously) agreed that dose to the bladder should be limited with a particular focus on limiting V_{30-35} Gy to prevent acute toxicity.

For other mandatory or recommended OAR (KQ3), data on dose response and dose volume relationships are scarce. The most well studied normal tissue, bone marrow, has inconsistent findings, hampering the establishment of clear optimization criteria (Supplement Table 1). For any other OAR outlined for an individual patient, dose should be reviewed as part of plan evaluation and prescription, and any patient specific considerations should be recorded.

The writing panel acknowledges that the definition of OAR dose constraints for IMRT treatment planning is highly dependent on clinical scenario and the acceptable risk for a giving patient/patient group. The UK RCR Rectal Cancer IMRT Guidance provides some guidance, primarily based on single institution experience [11].

KQ 7) Which dose metrics are to be considered for target dose evaluation?

IMRT plan optimisation for rectal cancer generally aims to deliver a homogenous dose prescription to the target volume(s); potentially with multiple different prescriptions to different volumes, as in the case of SIB delivery. Suggested planning objectives for target volumes were established through expert consensus, in line with general guidelines on IMRT planning and delivery [88], and informed by the UK Guidance [11]. Table 3 summarises the recommendations, using the recommended AAPM TG263 nomenclature [46]. In particular, note the suggested objectives for SIB delivery, ensuring evaluation of high dose conformity and minimising spillover of high dose into surrounding elective volumes.

KQ 8) How should dose to OAR and target volumes be prioritized during treatment plan optimization and evaluation?

Treatment plan optimisation and evaluation balances optimal target coverage with minimisation of dose to OAR, often necessitating compromises between the two objectives. The writing panel agreed (unanimous consensus) that the primary objective for standard preoperative radiotherapy for rectal cancer is to ensure adequate coverage of the GTV

Table 3
Target Volume Metrics, with standardised nomenclature according to American Association of Physicists in Medicine (AAPM) Task Group (TG) report 263.

Target volume	Standardised nomenclature (TG-263)	Dose metric	Objective†
Gross tumour, involved nodes, and clinical target volumes	GTvp, GTVn, CTVp, CTVn, CTVe*		Avoid cold spots in GTV / CTV target volumes; e.g. keep these volumes covered by 95% isodose. For plan evaluation, e.g. use $V_{95\%} \geq 99\%$
Planning target volume for primary tumour and involved lymph nodes (where relevant)	PTvp, PTVn	$D_{99\%}$ $D_{95\%}$ $D_{50\%}$	$\geq 90\%$ $\geq 95\%$ $= 100\% \pm 2\%$
Planning target volume for elective targets	PTVe*	$D_{2\%}$ $D_{99\%}$ $D_{95\%}$ $D_{50\%}$	$\leq 105\%$ $\geq 90\%$ $\geq 95\%$ $= 100\% \pm 2\%$
Evaluation structure when delivering simultaneously integrated boost (SIB)	PTVe minus (PTVp&n + 5 mm)		Avoid dose $\geq 105\%$ away from any SIB volumes Minimise volume receiving 107% of the elective volume prescription dose. For plan evaluation, e.g. use $V_{107\%} < 15\%$

† For dose measures (D_{xx}), all percentages are relative to the prescription dose for the volume in question.

* TG 263 does not have a recommended nomenclature for elective volumes. We propose using ‘CTVe’ and ‘PTVe’, or, alternatively, where there is no risk of ambiguity, simply ‘CTV’ and ‘PTV’.

Abbreviations: Gross Tumour Volume (GTV); Clinical Target Volume (CTV); Planning Target Volume (PTV); Dose to XX% of the volume ($D_{xx\%}$).

and CTV, to achieve effective tumour control with limited normal tissue exposure. The writing panel acknowledges that for individual patients and/or individual cases it will be at the treating physician discretion to prioritize an OAR over the target volume.

Whenever OAR constraints cannot be met due to unfavourable anatomy captured on the planning imaging, assessment should include if repeat planning imaging could improve the OAR constraints, see also KQ 2. The applied IGRT protocol should be in line with the planning prioritizations, see KQ14.

KQ 9) Are there any specific technical considerations to be made regarding IMRT planning?

The existing literature on IMRT planning for rectal cancer was reviewed, with a particular focus on papers reporting on technique comparisons, plan optimisation settings, and impact of contrast on optimisation. Key findings are summarised below, with specific tips and tricks based on the literature provided in Table 4 [25,89–105].

A number of studies have compared IMRT and VMAT planning and delivery techniques. In general, planning studies have found similar plan quality for IMRT compared to VMAT, with some inconsistencies in findings: For example, Shang et al. reported that OAR dose might be slightly better with IMRT [106], while Liu et al. [107] and Dunman et al. [108] found VMAT superior to IMRT with respect to target coverage and normal tissue sparing. Importantly, there is a clear and significant reduction in delivery time with VMAT compared to IMRT [92,106,108].

Recent literature, especially on newer treatment platforms, is sparse. Findings suggest that [109,110]. Studies reporting on helical tomotherapy planning and delivery indicate that it may be used to support normal tissue sparing [31], especially used in conduction with daily imaging [111].

It is worth noting that no studies were found with a direct comparison of plans from different treatment planning systems (TPS) with the same modality, treatment techniques, and planners. Consequently, it is

Table 4

Tips and tricks for optimisation of Intensity Modulated Radiotherapy (IMRT) and Volumetric Modulated Arc Therapy (VMAT), based on available literature.

IMRT optimisation	- Gantry angles avoiding anterior entry may be beneficial to keep OAR dose low (particularly bowel and bladder) [85] - Seven beam angles may provide a good compromise between dose conformality and complexity / delivery time [86], although fewer angles can also provide acceptable plans [87]
VMAT optimisation	- Dual arcs may allow one arc to focus on each lateral part of the target, sparing anterior organs at risk [88] - This may either be achieved by full dual arcs and avoidance volumes, or actively restricting the arcs to avoid anterior compartment [89] - A combination of an avoidance sector and a 'peritoneal space' avoidance volume might provide the best overall OAR sparing, at least for prone position [90] - Partial arcs give better and more robust dose distribution in the prone position, compared to full arcs [91] - Consider contracting the PTV used for optimisation away from the skin (see e.g. [92]) - Lower beam energies (6MV) are generally sufficient to achieve good plan quality [93], but higher beam energies could potentially be relevant for large patients / deep-seated tumours of large volumes [94]
Bowel	- Small bowel dose should be actively included in plan optimisation [23] - Bowel motion: - Although small bowel motion leads to uncertainties in delivered dose [95], IMRT plans still provide reduction in dose to small bowel compared to 3D-CRT plans [96] - Day-to-day bowel loop movements are significantly greater than day-to-day variation for peritoneal space [95]
Bladder	'Manually' optimised IMRT plans might not fully result in optimised bladder dose [97–98]; this might require additional attention if not utilising an auto-planning solution
Contrast and density corrections	- The presence of oral contrast on planning imaging may impact dose distributions, in particularly PTV and bowel dose [99], although with other literature indicating negligible impact (around 1% in mean dose to various structures) [100]. The overall impact likely depends on timing of CT, type of contrast, amount of contrast and dilution - Intravenous contrast agent has minimal impact on dose distributions [99] - Plans cannot be assumed to be robust to variation in intestinal gas, with particular risk to target coverage [101] - For plan optimisation, override densities that are likely to be different during delivery (oral contrast, significant rectal gas). Consider rescan for treatment planning when imaging includes significant gas which is unlikely to be present during treatment

Abbreviations: Organs at risk (OAR); Planning Target Volume (PTV); Computed Tomography (CT).

unclear what can be gained from improved optimisation engines alone. However, multicentre pre-trial benchmarking studies indicate that considerable variations can be found between centres, with different TPS and planners, even with stringent planning guidelines [102,112,113]. This supports the use of intercentre benchmarking and peer QA for treatment planning, including in the setting of clinical trials.

If SIB to the primary tumour is prescribed, IMRT or VMAT can deliver this without a substantial increase in dose to OAR [92,114–118], while achieving a more homogeneous dose distribution [119]. Importantly, when delivering a SIB, plan evaluation should include a measure of high dose conformity [120].

KQ 10) Can auto-planning solutions add value to IMRT planning, and what is the required QA?

Auto-planning is being increasingly utilised in radiotherapy, with solutions ranging from algorithm/scripting-based approaches to full AI based dose prediction and mimicking [121,122]. For IMRT planning for rectal cancer, auto-planning solutions contribute to improved OAR sparing [123–128], enhanced time efficiency [123–125,129], and increased consistency [98,129]. With one exception [130], all identified studies focused on treatment planning for LCRT. Auto-planning predominantly reduced the mean dose, V_{30Gy} and V_{40Gy} to the bladder and small bowel while maintaining comparable target coverage, indicating improved dose fall off towards these OAR.

Among the various automated planning approaches for rectal cancer, knowledge based planning (KBP) was the most widely utilized [97,98,124,126,128–130], followed by template-based iterative optimization [127] and script based optimization [123]. While these auto-planning solutions are commercially available, they do require some degree of in-house configuration and are not immediately deployable without customization. The latter two methods do not require any initial planning data to configure the auto-planning solution but the generation of a KBP model requires a minimum number of treatment plans as training data. Expanding the model library can improve the performance of the KBP model and increase its geometric diversity and representativeness. However, the introduction of suboptimal plans into the knowledge base should be avoided as this as this could negatively affect plan quality. Replanning suboptimal plans, either manually by an experienced dosimetrist or using a first iteration of the model, is beneficial to the performance of the model [126,129].

Although the reviewed studies demonstrated consistent findings, they were all retrospective and limited to single centre investigations, highlighting a lack of prospective implementation studies. Consequently, it remains uncertain whether the positive results found in the retrospective studies hold true in prospective clinical practice. It is recommended that each institution rigorously evaluates and monitors the performance of its auto-planning solution (whether in house or commercial) in prospective clinical practice so that the model can be updated if necessary [68].

KQ 11) What is the optimal method to calculate and report dose?

Modern TPS systems support type-B (e.g., convolution/superposition) and type-C (e.g., Monte Carlo and Grid Based Boltzmann solver) algorithms that enable dose calculations in a reasonable time frame. Consequently, the use of type-A dose calculation algorithms in the treatment planning of rectal cancer is not justified [131].

While there is a lack of definitive evidence supporting a single preferred method of dose reporting, consensus guidelines from the Global Quality Assurance of Radiation Therapy Clinical Trials Harmonisation Group [132] and NRG Clinical Trials [133] recommend dose to medium (D_m) as the standard framework for dose reporting.

Despite the fact that many dose constraints for these OAR have been established in terms of dose to water (D_w), it remains advisable to report in D_m to maintain consistency with ongoing and future clinical trials (expert consensus). Evaluation of D_w may allow comparison to historical dose prescriptions and tolerance thresholds (including those reviewed in KQ6). However, although it is technically feasible and supported by certain TPS to convert D_m to D_w (i.e., dose-to-water in medium, $D_{w,m}$), this practice is not recommended, as studies indicate that $D_{w,m}$ exhibits poorer concordance with D_w when compared to D_m [132,133]. In particular, the dose in cortical bone may be up to 10% higher when reported in $D_{w,m}$ compared to D_m , whereas the difference between D_m and D_w is of the order of a few percent, with D_m predicting a lower dose.

KQ 12) Is an MRI only workflow feasible using (commercially available) synthetic CT (sCT) for treatment planning?

Magnetic resonance imaging (MRI) is increasingly used in treatment planning for rectal cancer, due to the superior soft tissue contrast, especially for target definition [134]. Consequently, there is increasing interest in MRI only workflows to eliminate the need for treatment planning specific CT imaging and its registration uncertainties with MRI. The 2019 systematic review by Bird et al. found no clinical evidence

supporting the implementation of an MRI only workflow for rectal cancer [135], but our search identified seven studies published since this review, assessing the performance of commercially available MRI only software for rectal cancer [136–143]. In all studies, the standard MRI scanning protocol was supplemented with MRI sequences specific to the sCT generation software used. These sequences need to cover all the anatomy required for radiotherapy treatment planning, which sometimes requires two linked acquisitions to be stitched together offline into a single sequence to achieve sufficient superior-inferior scan length [136]. These sequences are optimised to minimise geometric distortions, and the geometric accuracy is routinely verified [137].

There is moderate evidence supporting the safe use of sCT for dose calculation in an MRI only workflow, with dose differences of less than 1% compared to planning CT across various imaging systems and commercially available sCT generation software [138,139,141–143]. However, some studies have reported a systematic overestimation of PTV dose [139,143] and general DVH differences [141]. These differences may be attributed to the handling of air pockets within the rectum by commercial software [139,143] or to the relative electron density curve that is applied to the generated sCT's HU values [141]. It is therefore recommended that the accuracy of a centre's local sCT generation technique is quantified during commissioning [137].

There is also moderate evidence to support the use of MRI or sCT as a reference image for position verification, demonstrating comparable results to CT, with systematic effects on Conebeam-CT (CBCT) registration below 1 mm and 0.5° [136,138–140,143]. However, commercial software currently lacks native support for using MRI or sCT as reference images, necessitating additional processing steps that can increase complexity [136,140]. Institutions should perform quality assurance on these processes to mitigate the associated risks.

In addition to improving the patient experience, patient comfort during MRI simulation should be optimised (expert consensus, unanimously) to minimize motion artifacts as well as minimizing inter- and intrafraction motion during treatment. This may be achieved using both earplugs and headphones, music for all patients, blankets to provide warmth, scanning of patients in the feet first position, and additional staff focus on providing information about the details of the MRI scan, particularly the scan time [144].

KQ 13) What are the setup and target volume shape uncertainties?

Understanding uncertainties in the treatment planning and delivery process is key to ensure the optimal benefit from IMRT and IGRT. Uncertainties may stem from a range of sources, including target definition and outlining, mechanical uncertainties in imaging and delivery systems, setup errors and inter- and intrafraction target shape variations. Following a comprehensive literature review, the focus of this key question was to examine setup variability and target volume shape variation. A thorough overview of the uncertainties is provided in a separate publication by Mantello et al [145].

Systematic and random errors in treatment setup, along with variations in target volume shape, were observed both between and within treatment fractions (interfraction and intrafraction) [34,35,146–172]. Despite the implementation of (online) IGRT, geometric variations in the CTV, particularly in the mesorectum, remained a significant and non-negligible source of interfraction and intrafraction uncertainty [35,147,148,150,151,153–159,162,163,166,168,170,172]. For the mesorectum, greater displacement was observed on the anterior surface of the upper half compared to the lower half; and conversely, the posterior surface exhibited more pronounced movement in the lower half compared to the upper half [34,147,148,150,151,162,163,169]. For the primary GTV, significant interfraction motion was also reported [148,150,155,158,166,170]. Position variation of individual involved lymph nodes appears to be strongly dependent on the position and specific lymph node compartment [173].

Non-negligible intrafractional organ motion was reported for both primary tumour and involved nodes, with additional increases observed for session durations exceeding 15 min [154,156,170].

As discussed in KQ 1, the use of a prone position with a bellyboard resulted in increased reported systematic and random setup errors [20,22,29,32].

In conclusion, systematic and random errors in setup and target volume shape variation are substantial and should be accounted for in the target volume margins. Setup errors can be effectively corrected using daily 2D or 3D IGRT, while target volume shape variations remain a challenge, emphasizing the importance of daily volumetric imaging (CBCT or MVCT) or adaptive workflows, to assess daily anatomical changes [174,175]. As different systematic and random errors were observed across the CTV sub-regions (GTVp, GTVn, mesorectum, and elective lymph node regions), the required margins should be defined separately (and possibly anisotropic) for each sub-region, reflecting their distinct motion patterns. For the sake of completeness, when calculating PTV margins, *all* errors in the treatment planning and delivery chain should be considered, not just those discussed here [174,176,177].

KQ 14) What is the optimal IGRT protocol to verify dose delivery?

An optimal IGRT protocol is closely associated with the expected interfractional and intrafractional positional variability of the target volume and surrounding OAR. A large number of studies have investigated interfractional and intrafractional uncertainties in patients with rectal cancer and especially interfraction target volume shape changes are large, see KQ13 and supporting paper Mantello et al [145]. Various factors have been identified that influence these changes, see KQ1 and KQ2. An additional critical factor is the choice of CTV to PTV margins and the underlying definition of the CTV. Based on these considerations, expert consensus (unanimously) is to always use daily volumetric imaging.

Decision making tools (e.g. traffic light protocols) should be defined (expert consensus, unanimously) in a multi-disciplinary setting to support (RTT-led) image evaluation and decision making in a daily volumetric imaging setting [11,174,178–180], see also Flowchart, Fig. 1 and the supporting images in Fig. 2. In these protocols, thresholds are typically defined for setup and anatomical changes, to ensure target coverage while minimizing dose to the surrounding healthy tissues.

Discussion

These ESTRO guidelines provide comprehensive recommendations for all technical aspects of radiotherapy planning and delivery with IMRT and IGRT for rectal cancer. With the widespread uptake of IMRT for rectal cancer [14], this guidance is expected to support broad clinical use.

However, in the process of synthesizing these guidelines, several key gaps in the literature became evident. When there was insufficient literature available, expert consensus was reached through voting. The panel was designed to be representative, including 2 RTTs, 2 MPEs, and 3 ROs. The members were a mix of academic and local experience and were geographically distributed across Europe. The small size of the group may have impacted on the outcome.

For imaging for treatment planning, it is currently unclear which strategies for bladder and rectal filling will optimally ensure consistent anatomy during treatment delivery. While most departments implement some form of patient preparation, high-quality data on this topic remains scarce [181].

For treatment planning, there is an overall lack of comprehensive dose volume response data for OAR in rectal cancer. This is particularly the case for modern treatment techniques (IMRT/VMAT) and for SCRT. Given the dependence of effective optimization of IMRT/VMAT treatment planning on this information [9], there is an obvious need for further research. Simply implementing IMRT may not be sufficient, if it is not clear which normal tissues to spare and how. This was illustrated by a recent paper from the RAPIDO trial, which compared toxicity in patients treated with and without IMRT, and found no benefit of the latter [9]. Despite the non-randomised nature of the comparison, and




Fig. 2.

the risk of centre-bias, these data provide a sobering reminder that newer techniques are not always a guarantee of improved outcomes. The evidence is even more lacking for optimisation of radiotherapy in emerging indications and settings, such as changing multimodality regimens like TNT or the combination with immuno- or targeted therapy. Dose-response and dose-volume effects may be impacted by such combination treatments. Other emerging scenarios (such as definitive radiotherapy for organ preservation) have seen almost no evaluation of normal tissue toxicity; and consequently, it is unclear which OAR should be delineated, optimised and evaluated in these settings [112,113]. Rectal wall, for instance, may potentially need to be considered as an OAR in the context of organ preservation where the rectum is not removed, but with no published data to date. There is a lack of data on normal tissue dose response for late toxicity for patients treated at higher dose levels (>55 Gy), especially where this has been part of an

organ preservation strategy. Additionally, there is a lack of patient-reported outcome data, with only a single study assessing patient-reported acute gastrointestinal symptoms [182].

A number of important topics were excluded from the current guideline from the outset. This included CTV and GTV definition: The former has already been covered by an ESTRO guideline [15], while the latter will be covered in a separate guideline under development by the ESTRO Lower GI guideline committee. As a direct consequence, the use of different imaging modalities for target volume definition, and other general radiological questions, such as MR sequencing for the MRI-only workflow (KQ12), were not covered here either. Also related to target volume definition, we deliberately choose not to provide simple recommendations for PTV margins, as optimal margins vary significantly depending on local departmental workflows. A comprehensive review of errors related to setup and target volume deformation can be found in a

separate publication and local margin calculations can be supported by these data [145].

Online volumetric imaging is strongly recommended, as it can identify patients where the margins used do not suffice and may require plan adaptation [11,176–180]. New tools have enabled other forms of adaptive radiotherapy [183,184 185–188]. The most promising technique is online adaptive radiotherapy [189–193]. This approach may be particularly advantageous for highly deformable targets, such as in rectal cancer, and is especially relevant in the context of dose escalation for organ preservation. Although promising in reducing treated volumes, these techniques were not included in this guideline due to limited accessibility and because clinical studies are still awaited to confirm this (e.g., NCT05883800, NCT06854679, NCT05108428).

Conclusion

We have provided guidelines for the clinical development and implementation of IMRT and IGRT. The overall level of evidence was often low and/or derived from expert consensus, underscoring the need for further research. Well-designed prospective clinical trials are essential to bridge existing knowledge gaps and enhance the quality of evidence regarding the application of IMRT and IGRT in the treatment of rectal cancer.

Disclaimer

ESTRO guidelines present scientific and medical opinions for educational and informational purposes only. Commercial use of any content in the guidelines without the prior written consent of ESTRO is strictly prohibited. The guidelines are not, and nor are they intended or implied to be, a substitute for professional medical advice or medical care. The advice of a medical professional should always be sought prior to commencing any form of medical treatment. Under no circumstances will ESTRO be held liable for any decision or action taken as a result of reliance on the content of the guidelines. ESTRO and all of its staff, agents and members disclaim any and all warranties and representations (implied or otherwise) with regards to the information contained on the guidelines. Each guideline is based on information available at the time the corresponding writing panel conducted its research and discussions on the topic. Guidelines should neither be deemed inclusive of all proper methods of care or factors influencing the decision making, nor exclusive of other methods reasonably directed to obtaining the same results. There may be new developments since publication that are not reflected in any one guideline and that may, over time, form the basis for ESTRO to revisit and update the guideline.

Funding Statement

The authors report no receipt of funds, grants or other support for the preparation of the manuscript.

CRediT authorship contribution statement

Rianne de Jong: Conceptualization, Data curation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review & editing. **Robin De Roover:** Conceptualization, Data curation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review & editing. **Lynsey Devlin:** Conceptualization, Data curation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review & editing. **Cihan Gani:** Conceptualization, Data curation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review & editing. **Daniel Portik:** Conceptualization, Data curation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review & editing. **Ane**

Appelt: Conceptualization, Data curation, Formal analysis, Methodology, Supervision, Visualization, Writing – original draft, Writing – review & editing.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Lynsey Devlin reports receipts of grants/research support with funding for research post provided by the Bearson Cancer Charity and by CRUK RadNet Glasgow. Ane Appelt reports receipt of grants/research support by research fellowship funded by Yorkshire Cancer Research and CRUK funding received for Leeds Radiotherapy Research Centre of Excellence. Daniel Portik reports receipts of grants/research support through research fellowship funded by EORTC Cancer Research Fund and Kom Op Tegen Kanker. Cihan Gani reports receipts of grants/research supports through financial and technical support for the department of Radiation Oncology Tubingen by Elekta and Philips.

Acknowledgments

The writing panel would like to thank the reviewing panel for their comprehensive review of these guidelines: Karin Haustermans, Corrie Marijnen, Jasper Nijkamp, Amanda Webster, Jackie Wu and the ASTRO Guideline subcommittee. We sincerely appreciate their valuable comments and suggestions, which helped improve the quality of the manuscript.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.radonc.2026.111409>.

References

- [1] Wo JY, et al. *Radiation Therapy for Rectal Cancer: an ASTRO Clinical Practice Guideline Focused Update*. *Pract Radiat Oncol* 2025;15:124–43.
- [2] Hofheinz RD, et al. *Localised rectal cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up*. *Ann Oncol* 2025;36:1007–24.
- [3] Dossa F, et al. *A watch-and-wait approach for locally advanced rectal cancer after a clinical complete response following neoadjuvant chemoradiation: a systematic review and meta-analysis*. *Lancet Gastroenterol Hepatol* 2017;2:501–13.
- [4] Slavu IM, et al. *A Review of Neoadjuvant Therapy and the Watch-and-wait Protocol in Rectal Cancer: current evidence and Future Directions*. *Cureus* 2024;16:e68461.
- [5] van der Valk MJM, et al. *Long-term outcomes of clinical complete responders after neoadjuvant treatment for rectal cancer in the International Watch & Wait Database (IWWD): an international multicentre registry study*. *Lancet* 2018;391:2537–45.
- [6] Bach SP, Collaborative S-T. *Can we save the rectum by watchful waiting or TransAnal surgery following (chemo)Radiotherapy versus Total mesorectal excision for early Rectal Cancer (STAR-TREC)? Protocol for the international, multicentre, rolling phase II/III partially randomized patient preference trial evaluating long-course concurrent chemoradiotherapy versus short-course radiotherapy organ preservation approaches*. *Colorectal Dis* 2022;24:639–51.
- [7] Peters FP, et al. *Mesorectal radiotherapy for early stage rectal cancer: a novel target volume*. *Clin Transl Radiat Oncol* 2020;21:104–11.
- [8] Wee CW, et al. *Intensity-modulated radiotherapy versus three-dimensional conformal radiotherapy in rectal cancer treated with neoadjuvant concurrent chemoradiation: a meta-analysis and pooled-analysis of acute toxicity*. *Jpn J Clin Oncol* 2018;48: 458–66.
- [9] Tanaka MD, et al. *Acute and late Radiation-Related Toxicity after Treatment of locally Advanced Rectal Cancer with Intensity Modulated Radiation Therapy Compared with 3-Dimensional Conformal Radiation Therapy in the RAPIDO Trial*. *Int J Radiat Oncol Biol Phys* 2025;123:470–82.
- [10] Appelt AL, et al. *Radiation dose-response model for locally advanced rectal cancer after preoperative chemoradiation therapy*. *Int J Radiat Oncol Biol Phys* 2013;85: 74–80.
- [11] Hanna CR, et al. *Intensity-modulated Radiotherapy for Rectal Cancer in the UK in 2020*. *Clin Oncol (R Coll Radiol)* 2021;33:214–23.
- [12] Teoh S, Muirhead R. *Rectal Radiotherapy–Intensity-modulated Radiotherapy delivery, Delineation and Doses Clin Oncol (R Coll Radiol)* 2016;28:93–102.
- [13] Vendrely V, et al. *Rectal cancer radiotherapy*. *Cancer Radiother* 2022;26:272–8.
- [14] Macnair A, et al. *Neoadjuvant Treatment of Rectal Cancer: a repeat UK-wide Survey after Implementation of National Intensity Modulated Radiotherapy Guidance*. *Clin Oncol (R Coll Radiol)* 2025;41:103835.

- [15] Valentini V, et al. *International consensus guidelines on Clinical Target volume delineation in rectal cancer*. *Radiother Oncol* 2016;120:195–201.
- [16] Das LJ, et al. *Efficacy of a belly board device with CT-simulation in reducing small bowel volume within pelvic irradiation fields*. *Int J Radiat Oncol Biol Phys* 1997;39:67–76.
- [17] Fu YT, Lam JC, Tze JM. *Measurement of irradiated small bowel volume in pelvic irradiation and the effect of a bellyboard*. *Clin Oncol (R Coll Radiol)* 1995;7:188–92.
- [18] Shanahan TG, et al. *Minimization of small bowel volume within treatment fields utilizing customized "belly boards"*. *Int J Radiat Oncol Biol Phys* 1990;19:469–76.
- [19] Hollenhorst H, et al. *Optimized radiation of pelvic volumes in the clinical setting by using a novel bellyboard with integrated gonadal shielding*. *Med Dosim* 2004;29:173–8.
- [20] Kim TH, et al. *Comparison of the belly board device method and the distended bladder method for reducing irradiated small bowel volumes in preoperative radiotherapy of rectal cancer patients*. *Int J Radiat Oncol Biol Phys* 2005;62:769–75.
- [21] Estabrook NC, et al. *Role of belly board device in the age of intensity modulated radiotherapy for pelvic irradiation*. *Med Dosim* 2016;41:300–4.
- [22] Hoffmann M, et al. *A critical literature review on the use of bellyboard devices to control small bowel dose for pelvic radiotherapy*. *Rep Pract Oncol Radiother* 2020;25:598–605.
- [23] Joye I, et al. *Implementation of volumetric modulated arc therapy for rectal cancer: pitfalls and challenges*. *Acta Oncol* 2015;54:1677–81.
- [24] Kim JY, et al. *Intensity-modulated radiotherapy with a belly board for rectal cancer*. *Int J Colorectal Dis* 2007;22:373–9.
- [25] Koeck J, et al. *Small bowel protection in IMRT for rectal cancer : a dosimetric study on supine vs. prone position*. *Strahlenther Onkol* 2017;193:578–88.
- [26] Nijkamp J, et al. *Bowel exposure in rectal cancer IMRT using prone, supine, or a belly board*. *Radiother Oncol* 2012;102:22–9.
- [27] Scobiala S, et al. *A treatment planning study of prone vs. supine positions for locally advanced rectal carcinoma : Comparison of 3-dimensional conformal radiotherapy, tomotherapy, volumetric modulated arc therapy, and intensity-modulated radiotherapy*. *Strahlenther Onkol* 2018;194:975–84.
- [28] Wiesendanger-Wittmer EM, et al. *Systematic review of the role of a belly board device in radiotherapy delivery in patients with pelvic malignancies*. *Radiother Oncol* 2012;102:325–34.
- [29] Yang Y, et al. *Effect of prone and supine treatment positions for postoperative treatment of rectal cancer on target dose coverage and small bowel sparing using intensity-modulated radiation therapy*. *Transl Cancer Res* 2020;9:491–9.
- [30] Kim A, et al. *The effect of prone and supine treatment positions for the pre-operative treatment of rectal cancer on organ-at-risk sparing and setup reproducibility using volumetric modulated arc therapy*. *Radiat Oncol* 2017;12:180.
- [31] Lin JC, et al. *Compared planning dosimetry of TOMO, VMAT and IMRT in rectal cancer with different simulated positions*. *Oncotarget* 2017;8:42020–9.
- [32] Siddiqui F, et al. *Image-guidance protocol comparison: supine and prone set-up accuracy for pelvic radiation therapy*. *Acta Oncol* 2008;47:1344–50.
- [33] Jong RH, W.; Betgen A et al. *Intra-fraction motion of patients in prone position compared to supine position*. *Radiother Oncol*, 2006. **128**(ESTRO 25): p. S49-S50.
- [34] Nijkamp J, et al. *Target volume shape variation during hypo-fractionated preoperative irradiation of rectal cancer patients*. *Radiother Oncol* 2009;92:202–9.
- [35] Nijkamp J, et al. *Target volume shape variation during irradiation of rectal cancer patients in supine position: comparison with prone position*. *Radiother Oncol* 2009;93:285–92.
- [36] Chang JS, et al. *Bladder filling variations during concurrent chemotherapy and pelvic radiotherapy in rectal cancer patients: early experience of bladder volume assessment using ultrasound scanner*. *Radiat Oncol J* 2013;31:41–7.
- [37] Yoon HI, et al. *Evaluating Variations of Bladder volume using an Ultrasound Scanner in Rectal Cancer patients During Chemoradiation: is Protocol-based Full Bladder Maintenance using a Bladder Scanner Useful to maintain the Bladder volume?* *PLoS One* 2015;10:e0128791.
- [38] Kim N, et al. *Simple calculation using anatomical features on pre-treatment verification CT for bladder volume estimation during radiation therapy for rectal cancer*. *BMC Cancer* 2020;20:942.
- [39] Feng X, et al. *Effectiveness of bladder filling control during online MR-guided adaptive radiotherapy for rectal cancer*. *Radiat Oncol* 2023;18:136.
- [40] Zhang H, et al. *Influence of bladder distension control on postoperative intensity-modulated radiotherapy in rectal cancer patients*. *Asian Pac J Cancer Prev* 2011;12:2969–72.
- [41] Potter R, et al. *The EMBRACE II study: the outcome and prospect of two decades of evolution within the GEC-ESTRO GYN working group and the EMBRACE studies*. *Clin Transl Radiat Oncol* 2018;9:48–60.
- [42] Gay HA, et al. *Pelvic normal tissue contouring guidelines for radiation therapy: a Radiation Therapy Oncology Group consensus panel atlas*. *Int J Radiat Oncol Biol Phys* 2012;83:e353–62.
- [43] Mir R, et al. *Organ at risk delineation for radiation therapy clinical trials: Global Harmonization Group consensus guidelines*. *Radiother Oncol* 2020;150:30–9.
- [44] Fuller CD, et al. *Prospective randomized double-blind pilot study of site-specific consensus atlas implementation for rectal cancer target volume delineation in the cooperative group setting*. *Int J Radiat Oncol Biol Phys* 2011;79:481–9.
- [45] Lin D, et al. *A Systematic Review of Contouring guidelines in Radiation Oncology: Analysis of Frequency, Methodology, and delivery of Consensus Recommendations*. *Int J Radiat Oncol Biol Phys* 2020;107:827–35.
- [46] Mayo CS, et al. *American Association of Physicists in Medicine Task Group 263: Standardizing Nomenclatures in Radiation Oncology*. *Int J Radiat Oncol Biol Phys* 2018;100:1057–66.
- [47] Brooks C, et al. *Proposed genitalia contouring guidelines in anal cancer intensity-modulated radiotherapy*. *Br J Radiol* 2015;88:20150032.
- [48] Arias F, et al. *Fecal incontinence and radiation dose on anal sphincter in patients with locally advanced rectal cancer (LARC) treated with preoperative chemoradiotherapy: a retrospective, single-institutional study*. *Clin Transl Oncol* 2017;19:969–75.
- [49] Yi SK, et al. *Development of a standardized method for contouring the lumbosacral plexus: a preliminary dosimetric analysis of this organ at risk among 15 patients treated with intensity-modulated radiotherapy for lower gastrointestinal cancers and the incidence of radiation-induced lumbosacral plexopathy*. *Int J Radiat Oncol Biol Phys* 2012;84:376–82.
- [50] Augurio A, et al. *Contouring of emerging organs-at-risk (OARS) of the female pelvis and interobserver variability: a study by the Italian association of radiotherapy and clinical oncology (AIRO)*. *Clin Transl Radiat Oncol* 2023;43:100688.
- [51] Kronborg CJS, et al. *Defining and sparing sexual function-related organs at risk for rectal cancer radiotherapy*. *Acta Oncol* 2025;64:1136–42.
- [52] Harrison K, et al. *Machine Learning for Auto-Segmentation in Radiotherapy Planning*. *Clin Oncol* 2022;34:74–88.
- [53] Brouwer CL, et al. *Machine learning applications in radiation oncology: current use and needs to support clinical implementation*. *Physics & Imaging in Radiation Oncology* 2020;16:144–8.
- [54] Li J, et al. *Clinical evaluation on automatic segmentation results of convolutional neural networks in rectal cancer radiotherapy*. *Front Oncol* 2023;13:1158315.
- [55] Sha X, et al. *Clinical target volume and organs at risk segmentation for rectal cancer radiotherapy using the Flex U-Net network*. *Front Oncol* 2023;13:1172424.
- [56] Sartor H, et al. *Auto-segmentations by convolutional neural network in cervical and anorectal cancer with clinical structure sets as the ground truth*. *Clin Transl Radiat Oncol* 2020;25:37–45.
- [57] Men K, Dai J, Li Y. *Automatic segmentation of the clinical target volume and organs at risk in the planning CT for rectal cancer using deep dilated convolutional neural networks*. *Med Phys* 2017;44:6377–89.
- [58] Gambacorta MA, et al. *Clinical validation of atlas-based auto-segmentation of pelvic volumes and normal tissue in rectal tumors using auto-segmentation computed system*. *Acta Oncol* 2013;52:1676–81.
- [59] Men K, et al. *A study of positioning orientation effect on segmentation accuracy using convolutional neural networks for rectal cancer*. *J Appl Clin Med Phys* 2019;20:110–7.
- [60] Mackay K, et al. *A Review of the Metrics used to Assess Auto-Contouring Systems in Radiotherapy*. *Clin Oncol (R Coll Radiol)* 2023;35:354–69.
- [61] Radici L, et al. *Implementation of a Commercial Deep Learning-based Auto Segmentation Software in Radiotherapy: Evaluation of Effectiveness and Impact on Workflow*. *Life (Basel)* 2022;12.
- [62] Lempart M, et al. *Pelvic U-Net: multi-label semantic segmentation of pelvic organs at risk for radiation therapy and anal cancer patients using a deeply supervised shuffle attention convolutional neural network*. *Radiat Oncol* 2022;17:114.
- [63] Zhou H, et al. *A deep learning based automatic segmentation approach for anatomical structures in intensity modulation radiotherapy*. *Math Biosci Eng* 2021;18:7506–24.
- [64] van Heeswijk MM, et al. *Automated and Semiautomated Segmentation of Rectal Tumor Volumes on Diffusion-Weighted MRI: can it Replace Manual Volumetry?* *Int J Radiat Oncol Biol Phys* 2016;94:824–31.
- [65] Strolin S, et al. *How smart is artificial intelligence in organs delineation? Testing a CE and FDA-approved Deep-Learning tool using multiple expert contours delineated on planning CT images*. *Front Oncol* 2023;13:1089807.
- [66] Wu Y, et al. *A blind randomized validated convolutional neural network for auto-segmentation of clinical target volume in rectal cancer patients receiving neoadjuvant radiotherapy*. *Cancer Med* 2022;11:166–75.
- [67] Gambacorta MA, et al. *Automatic segmentation software in locally advanced rectal cancer: READY (REsearch program in Auto Delineation sYstem)-RECTAL 02: prospective study*. *Oncotarget* 2016;7:42579–84.
- [68] Vandewinckele L, et al. *Overview of artificial intelligence-based applications in radiotherapy: Recommendations for implementation and quality assurance*. *Radiother Oncol* 2020;153:55–66.
- [69] Hou Z, et al. *Clinical evaluation of deep learning-based automatic clinical target volume segmentation: a single-institution multi-site tumor experience*. *Radiol Med* 2023;128:1250–61.
- [70] Knuth F, et al. *Semi-automatic tumor segmentation of rectal cancer based on functional magnetic resonance imaging*. *Phys Imaging Radiat Oncol* 2022;22:77–84.
- [71] Trebeschi S, et al. *Deep Learning for Fully-Automated Localization and Segmentation of Rectal Cancer on Multiparametric MR*. *Sci Rep* 2017;7:5301.
- [72] Geng J, et al. *Towards deep-learning (DL) based fully automated target delineation for rectal cancer neoadjuvant radiotherapy using a divide-and-conquer strategy: a study with multicenter blind and randomized validation*. *Radiat Oncol* 2023;18:164.
- [73] Men K, et al. *Automatic segmentation of three clinical target volumes in radiotherapy using lifelong learning*. *Radiother Oncol* 2021;157:1–7.
- [74] Larsson R, et al. *Automatic Delineation of the Clinical Target volume in Rectal Cancer for Radiation Therapy using Three-dimensional fully Convolutional Neural Networks*. *Annu Int Conf IEEE Eng Med Biol Soc* 2018;2018:5898–901.
- [75] Guo H, et al. *The dosimetric impact of deep learning-based auto-segmentation of organs at risk on nasopharyngeal and rectal cancer*. *Radiat Oncol* 2021;16:113.
- [76] Ciernik IF, et al. *Automated functional image-guided radiation treatment planning for rectal cancer*. *Int J Radiat Oncol Biol Phys* 2005;62:893–900.
- [77] Anders LC, et al. *Performance of an atlas-based autosegmentation software for delineation of target volumes for radiotherapy of breast and anorectal cancer*. *Radiother Oncol* 2012;102:68–73.
- [78] Holyoake DLP, Partridge M, Hawkins MA. *Systematic review and meta-analysis of small bowel dose-volume and acute toxicity in conventionally-fractionated rectal cancer radiotherapy*. *Radiother Oncol* 2019;138:38–44.

- [79] Ellsworth SG, et al. *Declarations of Independence: how embedded Multicollinearity Errors Affect Dosimetric and other complex analyses in Radiation Oncology*. Int J Radiat Oncol Biol Phys 2023;117:1054–62.
- [80] Yang TJ, et al. *Predictors of acute gastrointestinal toxicity during pelvic chemoradiotherapy in patients with rectal cancer*. Gastrointest Cancer Res 2013;6:129–36.
- [81] Banerjee R, et al. *Small bowel dose parameters predicting grade ≥ 3 acute toxicity in rectal cancer patients treated with neoadjuvant chemoradiation: an independent validation study comparing peritoneal space versus small bowel loop contouring techniques*. Int J Radiat Oncol Biol Phys 2013;85:1225–31.
- [82] Myerson RJ, et al. *A phase I/II trial of three-dimensionally planned concurrent boost radiotherapy and protracted venous infusion of 5-FU chemotherapy for locally advanced rectal carcinoma*. Int J Radiat Oncol Biol Phys 2001;50:1299–308.
- [83] Abraham AG, et al. *Dosimetric Parameters predicting late Small Bowel Toxicity in patients with Rectal Cancer Receiving Neoadjuvant Chemoradiation*. Pract Radiat Oncol 2021;11:e70–9.
- [84] Chopra S, et al. *Predictors of late bowel toxicity using three different methods of contouring in patients undergoing post-operative radiation for cervical cancer*. Br J Radiol 2015;88:20150054.
- [85] Hysing LB, et al. *Planning organ at risk volume margins for organ motion of the intestine*. Radiother Oncol 2006;80:349–54.
- [86] Dominello MM, et al. *Limitations of the bowel bag contouring technique in the definitive treatment of cervical cancer*. Pract Radiat Oncol 2014;4:e15–20.
- [87] Appelt AL, et al. *Dose-response of acute urinary toxicity of long-course preoperative chemoradiotherapy for rectal cancer*. Acta Oncol 2015;54:179–86.
- [88] Hodapp N. *The ICRU Report 83: prescribing, recording and reporting photon-beam intensity-modulated radiation therapy (IMRT)*. Strahlenther Onkol 2012;188:97–9.
- [89] Duthoy W, et al. *Clinical implementation of intensity-modulated arc therapy (IMAT) for rectal cancer*. Int J Radiat Oncol Biol Phys 2004;60:794–806.
- [90] Mok H, et al. *Intensity modulated radiation therapy (IMRT): differences in target volumes and improvement in clinically relevant doses to small bowel in rectal carcinoma*. Radiat Oncol 2011;6:63.
- [91] Guerrero Urbano MT, et al. *Intensity-modulated radiotherapy in patients with locally advanced rectal cancer reduces volume of bowel treated to high dose levels*. Int J Radiat Oncol Biol Phys 2006;65:907–16.
- [92] Cilla S, et al. *Volumetric modulated arc therapy with simultaneous integrated boost for locally advanced rectal cancer*. Clin Oncol (R Coll Radiol) 2012;24:261–8.
- [93] Wang CC, et al. *Neoadjuvant bevacizumab and chemoradiotherapy in locally advanced rectal cancer: early outcome and technical impact on toxicity*. World J Surg Oncol 2014;12:329.
- [94] Brennan VS, et al. *A novel dynamic arc treatment planning solution to reduce dose to small bowel in preoperative radiotherapy for rectal cancer*. Med Dosim 2019;44:258–65.
- [95] Zhang Z, et al. *Advantages and robustness of partial VMAT with prone position for neoadjuvant rectal cancer evaluated by CBCT-based offline adaptive radiotherapy*. Radiat Oncol 2023;18:102.
- [96] Huang MY, et al. *Helical tomotherapy combined with capecitabine in the preoperative treatment of locally advanced rectal cancer*. Biomed Res Int 2014;2014:352083.
- [97] Huang Y, et al. *Impact of nominal photon energies on normal tissue sparing in knowledge-based radiotherapy treatment planning for rectal cancer patients*. PLoS One 2019;14:e0213271.
- [98] Hirashima H, et al. *Reducing variability among treatment machines using knowledge-based planning for head and neck, pancreatic, and rectal cancer*. J Appl Clin Med Phys 2021;22:245–54.
- [99] Li S, et al. *Evaluation of small bowel motion and feasibility of using the peritoneal space to replace bowel loops for dose constraints during intensity-modulated radiotherapy for rectal cancer*. Radiat Oncol 2020;15:211.
- [100] Nuytens JJ, et al. *The influence of small bowel motion on both a conventional three-field and intensity modulated radiation therapy (IMRT) for rectal cancer*. Cancer Radiother 2004;8:297–304.
- [101] Vaniqui A, et al. *Treatment plan quality assessment for radiotherapy of rectal cancer patients using prediction of organ-at-risk dose metrics*. Phys Imaging Radiat Oncol 2020;16:74–80.
- [102] van den Ende RPJ, et al. *Radiotherapy quality assurance for mesorectum treatment planning within the multi-center phase II STAR-TRC trial: dutch results*. Radiat Oncol 2020;15:41.
- [103] Jing H, et al. *Oral contrast agents lead to underestimation of dose calculation in volumetric-modulated arc therapy planning for pelvic irradiation*. Chin Med J (Engl) 2020;133:2061–70.
- [104] Joseph K, et al. *Dosimetric effect of small bowel oral contrast on conventional radiation therapy, linear accelerator-based intensity modulated radiation therapy, and helical tomotherapy plans for rectal cancer*. Pract Radiat Oncol 2015;5:e95–.
- [105] Radu C, et al. *Integrated peripheral boost in preoperative radiotherapy for the locally most advanced non-resectable rectal cancer patients*. Acta Oncol 2013;52:528–37.
- [106] Shang J, et al. *VMAT planning study in rectal cancer patients*. Radiat Oncol 2014;9:219.
- [107] Liu M, et al. *Dosimetric comparative study of 3 different postoperative radiotherapy techniques (3D-CRT, IMRT, and RapidArc) for II-III stage rectal cancer*. Medicine (Baltimore) 2015;94:e372.
- [108] Duman E, Bilek Y, Ceyran G. *A comparison of radiotherapy treatment planning techniques in patients with rectal cancers by analyzing testes doses*. J Cancer Res Ther 2021;17:56–61.
- [109] Zheng J, Xia Y, Sun L. *A Comprehensive Evaluation of the Application of the Halcyon (2.0) IMRT Technique in Long-Course Radiotherapy for Rectal Cancer*. Technol Cancer Res Treat 2022;21:15330338221074501.
- [110] Sakanaka K, et al. *Preoperative chemoradiotherapy for locally advanced low rectal cancer using intensity-modulated radiotherapy to spare the intestines: a single-institutional pilot trial*. J Radiat Res 2022;63:88–97.
- [111] Engels B, et al. *Phase II study of preoperative helical tomotherapy with a simultaneous integrated boost for rectal cancer*. Int J Radiat Oncol Biol Phys 2012;83:142–8.
- [112] Appelt AL, et al. *Robust dose planning objectives for mesorectal radiotherapy of early stage rectal cancer - a multicentre dose planning study*. Tech Innov Patient Support Radiat Oncol 2019;11:14–21.
- [113] Arp DT, et al. *Treatment planning for patients with low rectal cancer in a multicenter prospective organ preservation study*. Phys Med 2024;118:103206.
- [114] Seierstad T, et al. *MR-guided simultaneous integrated boost in preoperative radiotherapy of locally advanced rectal cancer following neoadjuvant chemotherapy*. Radiother Oncol 2009;93:279–84.
- [115] Li JL, et al. *Preoperative concomitant boost intensity-modulated radiotherapy with oral capecitabine in locally advanced mid-low rectal cancer: a phase II trial*. Radiother Oncol 2012;102:4–9.
- [116] Bakkal BH, Elmas O. *Dosimetric comparison of organs at risk in 5 different radiotherapy plans in patients with preoperatively irradiated rectal cancer*. Medicine (Baltimore) 2021;100:e24266.
- [117] Owens R, et al. *Intensity-Modulated Radiotherapy with a Simultaneous Integrated boost in Rectal Cancer*. Clin Oncol (R Coll Radiol) 2020;32:35–42.
- [118] Jankarashvili N, et al. *Neoadjuvant volumetric modulated arc radiochemotherapy with a simultaneous integrated boost technique compared to standard chemoradiation for locally advanced rectal cancer*. Turk J Med Sci 2019;49:1484–9.
- [119] Karaca S, Aysenur Arli Karacam K. *A comparison of dosimetric and clinical parameters between different IMRT boost techniques in preoperative rectal cancer*. J BUON, 2021;26(4):231–1238.
- [120] Yang Y, et al. *A Dosimetric Analysis of Preoperative Intensity-modulated and Image-guided Radiation Therapy with and without Simultaneous Integrated boost for locally Advanced Rectal Cancer*. Technol Cancer Res Treat 2015;14:557–63.
- [121] Hussein M, et al. *Automation in intensity modulated radiotherapy treatment planning - a review of recent innovations*. Br J Radiol 2018;91:20180270.
- [122] Nguyen D, et al. *Advances in Automated Treatment Planning*. Semin Radiat Oncol 2022;32:343–50.
- [123] Hirotsaki K, et al. *Fully automated volumetric modulated arc therapy planning for locally advanced rectal cancer: feasibility and efficiency*. Radiat Oncol 2023;18:147.
- [124] Shepherd M, et al. *Developing knowledge-based planning for gynaecological and rectal cancers: a clinical validation of RapidPlan ()*. J Med Radiat Sci 2020;67:217–24.
- [125] Song Y, et al. *Fully automatic volumetric modulated arc therapy plan generation for rectal cancer*. Radiother Oncol 2016;119:531–6.
- [126] Wang M, et al. *An interactive plan and model evolution method for knowledge-based pelvic VMAT planning*. J Appl Clin Med Phys 2018;19:491–8.
- [127] Wortel G, et al. *Characterization of automatic treatment planning approaches in radiotherapy*. Phys Imaging Radiat Oncol 2021;19:60–5.
- [128] Wu H, et al. *Applying a RapidPlan model trained on a technique and orientation to another: a feasibility and dosimetric evaluation*. Radiat Oncol 2016;11:108.
- [129] Wu H, et al. *A dosimetric evaluation of knowledge-based VMAT planning with simultaneous integrated boosting for rectal cancer patients*. J Appl Clin Med Phys 2016;17:78–85.
- [130] Huang K, et al. *Automatic end-to-end VMAT treatment planning for rectal cancers*. J Appl Clin Med Phys 2024;25:e14259.
- [131] Martino D. *Dose Calculation Algorithms for External Radiation Therapy: an Overview for Practitioners*. Appl Sci 2021;11.
- [132] Kry SF, et al. *Report dose-to-medium in clinical trials where available; a consensus from the Global Harmonisation Group to maximize consistency*. Radiother Oncol 2021;159:106–11.
- [133] Gladstone DJ, et al. *Dose Specification for NRG Radiation Therapy Trials*. Int J Radiat Oncol Biol Phys 2016;95:1344–5.
- [134] Horvat N, et al. *MRI of Rectal Cancer: Tumor Staging, Imaging Techniques, and Management*. Radiographics 2019;39:367–87.
- [135] Bird D, et al. *A Systematic Review of the Clinical Implementation of Pelvic magnetic Resonance Imaging-only Planning for External Beam Radiation Therapy*. Int J Radiat Oncol Biol Phys 2019;105:479–92.
- [136] Bird D, et al. *Patient position verification in magnetic-resonance imaging only radiotherapy of anal and rectal cancers*. Phys Imaging Radiat Oncol 2021;19:72–7.
- [137] Glide-Hurst CK, et al. *Task group 284 report: magnetic resonance imaging simulation in radiotherapy: considerations for clinical implementation, optimization, and quality assurance*. Med Phys 2021;48:e636–70.
- [138] Kemppainen R, et al. *Assessment of dosimetric and positioning accuracy of a magnetic resonance imaging-only solution for external beam radiotherapy of pelvic anatomy*. Phys Imaging Radiat Oncol 2019;11:1–8.
- [139] Maspero M, et al. *Feasibility of magnetic resonance imaging-only rectum radiotherapy with a commercial synthetic computed tomography generation solution*. Phys Imaging Radiat Oncol 2018;7:58–64.
- [140] O'Connor LM, et al. *Cone beam computed tomography image guidance within a magnetic resonance imaging-only planning workflow*. Phys Imaging Radiat Oncol 2023;27:100472.
- [141] O'Connor LM, et al. *Optimisation and validation of an integrated magnetic resonance imaging-only radiotherapy planning solution*. Phys Imaging Radiat Oncol 2021;20:34–9.
- [142] Wyatt JJ, et al. *Comprehensive dose evaluation of a Deep Learning based synthetic Computed Tomography algorithm for pelvic magnetic Resonance-only radiotherapy*. Radiother Oncol 2023;184:109692.
- [143] Yu VY, et al. *A multi-institutional analysis of a general pelvis continuous Hounsfield unit synthetic CT software for radiotherapy*. J Appl Clin Med Phys 2021;22:207–15.

- [144] Bird DP, S., Gilbert A, Casanova N, Cooper R et al. Assessing the patient experience of anal and rectal cancer MR simulation for radiotherapy treatment planning. *J Radiother Pract*, 2022;21:487–92.
- [145] Mantello G. *Setup and target volume shape variation in rectal cancer radiotherapy: a systematic literature review*. Tech Innovat Patient Support Radiat Oncol 2025.
- [146] Bansal S, et al. Determination of Optimal Clinical Target volume to Planning Target volume Margins for Conformal Radiotherapy Planning using image Guidance System in Rectal Cancer in Prone Position. *J Med Phys* 2019;44:65–7.
- [147] Boldrini L, et al. Mesorectal motion evaluation in rectal cancer MR-guided radiotherapy: an exploratory study to quantify treatment margins. *Radiat Oncol* 2023;18:4.
- [148] Brierley JD, et al. Rectal motion in patients receiving preoperative radiotherapy for carcinoma of the rectum. *Int J Radiat Oncol Biol Phys* 2011;80:97–102.
- [149] Chen YJ, et al. Setup variations in radiotherapy of anal cancer: advantages of target volume reduction using image-guided radiation treatment. *Int J Radiat Oncol Biol Phys* 2012;84:289–95.
- [150] Chong I, et al. Quantification of organ motion during chemoradiotherapy of rectal cancer using cone-beam computed tomography. *Int J Radiat Oncol Biol Phys* 2011;81:e431–8.
- [151] Cox MC, et al. Mesorectal shape variation in rectal cancer radiotherapy in prone position using a belly board. *Phys Imaging Radiat Oncol* 2021;19:120–5.
- [152] Cubillos Mesias M, et al. Quantification and Assessment of Interfraction Setup Errors based on Cone Beam CT and Determination of Safety Margins for Radiotherapy. *PLoS One* 2016;11:e0150326.
- [153] Durrant L, et al. Quantifying target-specific motion in anal cancer patients treated with intensity modulated radiotherapy (IMRT). *Radiother Oncol* 2016;121:92–7.
- [154] Eijkelkamp H, et al. Planning target volume margin assessment for online adaptive MR-guided dose-escalation in rectal cancer on a 1.5 T MR-Linac. *Radiother Oncol* 2021;162:150–5.
- [155] Garibaldi C, et al. Cone-beam CT-based inter-fraction localization errors for tumors in the pelvic region. *Phys Med* 2018;46:59–66.
- [156] Kensen CM, et al. Online Adaptive MRI-Guided Radiotherapy for Primary Tumor and Lymph Node Boosting in Rectal Cancer. *Cancers (Basel)* 2023;15.
- [157] Kershaw L, et al. Image Guided Radiation Therapy strategies for Pelvic Lymph Node Irradiation in High-Risk Prostate Cancer: Motion and Margins. *Int J Radiat Oncol Biol Phys* 2018;100:68–77.
- [158] Kleijnen JJE, et al. MRI-based tumor inter-fraction motion statistics for rectal cancer boost radiotherapy. *Acta Oncol* 2019;58:232–6.
- [159] Marnouche EA, et al. Evaluation of margins in pelvic lymph nodes and prostate radiotherapy and the impact of bladder and rectum on prostate position. *Cancer Radiother* 2021;25:161–8.
- [160] Miyamoto J, et al. The Role of Image-guided Radiotherapy in the Treatment of Anorectal Cancer using Prone Belly-board Positioning. *Anticancer Res* 2016;36:3013–7.
- [161] Mohamed R, et al. Factors Affecting Isocenter Displacement and Planning Target volume Margin for patients with Rectal Cancer Receiving Radiation Therapy. *Adv Radiat Oncol* 2022;7:101060.
- [162] Moldovan N, et al. Evaluation of the ITV-margin and variables affecting bladder and mesorectal deformation during long course neoadjuvant radiotherapy for rectal cancer. *Med Dosim* 2022;47:236–41.
- [163] Muren LP, et al. On the use of margins for geometrical uncertainties around the rectum in radiotherapy planning. *Radiother Oncol* 2004;70:11–9.
- [164] Patni N, et al. Assessment of three-dimensional setup errors in image-guided pelvic radiotherapy for uterine and cervical cancer using kilovoltage cone-beam computed tomography and its effect on planning target volume margins. *J Cancer Res Ther* 2017;13:131–6.
- [165] Rajeev KR, et al. A comparative study of set up variations and bowel volumes in supine versus prone positions of patients treated with external beam radiation for carcinoma rectum. *J Cancer Res Ther* 2014;10:937–41.
- [166] Raso R, et al. Assessment and clinical validation of margins for adaptive simultaneous integrated boost in neo-adjuvant radiochemotherapy for rectal cancer. *Phys Med* 2015;31:167–72.
- [167] Robert N, et al. Rotational Set up Uncertainty in Non-6D Couch and its Effects in Clinical Target Volume- Planning Target volume Margin Calculation for Different Sites. *J Med Phys* 2023;48:43–9.
- [168] Schippers MG, et al. Position shifts and volume changes of pelvic and para-aortic nodes during IMRT for patients with cervical cancer. *Radiother Oncol* 2014;111:442–5.
- [169] Tournel K, et al. Assessment of intrafractional movement and internal motion in radiotherapy of rectal cancer using megavoltage computed tomography. *Int J Radiat Oncol Biol Phys* 2008;71:934–9.
- [170] van den Ende RPJ, et al. Feasibility of Gold Fiducial Markers as a Surrogate for Gross Tumor volume Position in Image-Guided Radiation Therapy of Rectal Cancer. *Int J Radiat Oncol Biol Phys* 2019;105:1151–9.
- [171] Wu WC, et al. Impact of body-mass factors on setup displacement during pelvic irradiation in patients with lower abdominal cancer. *Radiol Oncol* 2019;53:256–64.
- [172] Nijkamp J, et al. Repeat CT assessed CTV variation and PTV margins for short- and long-course pre-operative RT of rectal cancer. *Radiother Oncol* 2012;102:399–405.
- [173] Arp DT, et al. Individual lymph node position variation for rectal cancer patients treated with long course chemoradiotherapy. *Phys Imaging Radiat Oncol* 2024;31:100599.
- [174] McNair HA, Franks KN, van Herk M. On Target 2: Updated Guidance for Image-guided Radiotherapy. *Clin Oncol (R Coll Radiol)* 2022;34:187–8.
- [175] Tudor. *Geometric Uncertainties in Daily Online IGRT: refining the CTV-PTV Margin for Contemporary Radiotherapy*. Brit Inst Radiol 2020.
- [176] van Herk M. Errors and margins in radiotherapy. *Semin Radiat Oncol* 2004;14:52–64.
- [177] van Herk M, et al. The probability of correct target dosage: dose-population histograms for deriving treatment margins in radiotherapy. *Int J Radiat Oncol Biol Phys* 2000;47:1121–35.
- [178] Alexander SE, et al. RTT-led IGRT for cervix cancer; training, implementation and validation. *Tech Innov Patient Support Radiat Oncol* 2019;12:41–9.
- [179] Buijs M, et al. Take Action Protocol: a radiation therapist led approach to act on anatomical changes seen on CBCT. *Tech Innov Patient Support Radiat Oncol* 2021;17:71–7.
- [180] Hattu D, et al. A traffic light protocol workflow for image-guided adaptive radiotherapy in lung cancer patients. *Radiother Oncol* 2022;175:152–8.
- [181] Slevin F. Overview of patient preparation strategies to manage internal organ motion during radiotherapy in the pelvis. *J Radiother Pract* 2020;19:182–9.
- [182] Chen RC, et al. Dose-volume effects on patient-reported acute gastrointestinal symptoms during chemoradiation therapy for rectal cancer. *Int J Radiat Oncol Biol Phys* 2012;83:e513–7.
- [183] Sonke JJ, Aznar M, Rasch C. Adaptive Radiotherapy for Anatomical changes. *Semin Radiat Oncol* 2019;29:245–57.
- [184] Nijkamp J, et al. Adaptive radiotherapy for long course neo-adjuvant treatment of rectal cancer. *Radiother Oncol* 2012;103:353–9.
- [185] Chen D, et al. Efficient strategy for magnetic resonance image-guided adaptive radiotherapy of rectal cancer using a library of reference plans. *Phys Imaging Radiat Oncol* 2025;33:100747.
- [186] de Jong R, et al. Dosimetric benefit of an adaptive treatment by means of plan selection for rectal cancer patients in both short and long course radiation therapy. *Radiat Oncol* 2020;15:13.
- [187] Lutkenhaus LJ, et al. Potential dosimetric benefit of an adaptive plan selection strategy for short-course radiotherapy in rectal cancer patients. *Radiother Oncol* 2016;119:525–30.
- [188] Beekman C, et al. Margin and PTV volume reduction using a population based library of plans strategy for rectal cancer radiotherapy. *Med Phys* 2018;45:4345–54.
- [189] Schuurhuizen C, et al. Clinical feasibility of treatment planning on a diagnostic CT scan without or with single fraction plan adaptation in patients with stage II/III rectal cancer. *Radiother Oncol* 2025;206:110840.
- [190] de Jong R, et al. Feasibility of Conebeam CT-based online adaptive radiotherapy for neoadjuvant treatment of rectal cancer. *Radiat Oncol* 2021;16:136.
- [191] Kensen CM, et al. Effect of intrafraction adaptation on PTV margins for MRI guided online adaptive radiotherapy for rectal cancer. *Radiat Oncol* 2022;17:110.
- [192] Intven MPW, et al. Online adaptive MR-guided radiotherapy for rectal cancer; feasibility of the workflow on a 1.5T MR-linac: clinical implementation and initial experience. *Radiother Oncol* 2021;154:172–8.
- [193] Henning J, et al. CT-Based Adaptive Short Course Radiotherapy for Rectal Cancer: a Case Report. *Cureus* 2024;16:e69264.

Rianne de Jong^{a,*}, Robin De Roover^b, Lynsey Devlin^{c,d}, Cihan Gani^e, Giovanna Mantello^f, Daniel Portik^{g,h}, Ane Appelt^{i,j}

^a Department of Radiation Oncology, Leiden University Medical Centre, Leiden, the Netherlands

^b Department of Radiation Oncology, University Hospitals Leuven, Leuven, Belgium

^c Department of Radiotherapy, The Beatson West of Scotland Cancer Centre, Glasgow, United Kingdom

^d CRUK RadNet Glasgow, University of Glasgow, Glasgow, United Kingdom

^e Department of Radiation Oncology, University Hospital and Medical Faculty Tübingen, Eberhard Karls University Tübingen, Tübingen, Germany

^f Radiotherapy Department, Azienda Ospedaliero Universitaria della Marche, 60126 Ancona, Italy

^g Department of Radiation Oncology (Mastro), GROW Research Institute for Oncology and Reproduction, Maastricht University Medical Centre+, Maastricht, the Netherlands

^h European Organisation for Research and Treatment of Cancer (EORTC) Headquarters, Brussels, Belgium

ⁱ Leeds Institute of Medical Research, University of Leeds, Leeds, United Kingdom

^j Department of Oncology, Section of Radiotherapy, Rigshospitalet - Technical University Hospital of Greater Copenhagen, Copenhagen, Denmark

* Corresponding author at: Department of Radiation Oncology, Leiden University Medical Centre, Postbus 9600 | 2300 RC Leiden. E-mail address: m.a.j.de.jong@lumc.nl (R. de Jong).